

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 2 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

RECEIVED
CENTRAL FAX CENTER

JUL 18 2005

In the Claims

Please amend the claims by replacing all prior versions, and listings, of claims pursuant to 37 C.F.R. §1.121(c) as follows:

1-122. (Canceled)

123. (Currently Amended) In a process for obtaining a pharmaceutical product containing a mixture of polypeptides, each of which consists essentially of alanine, glutamic acid, tyrosine and lysine, wherein the mixture has an average molecular weight from 4000 to 13,000 Daltons and in the mixture the molar fraction of alanine is 0.427, of glutamic acid is 0.141, of lysine is 0.337 and of tyrosine is 0.093 and wherein during the process includes determining the molecular weight distribution of a batch of a-an aqueous mixture of polypeptides, each of which consists essentially of alanine, glutamic acid, tyrosine and lysine, is tested using a gel permeation chromatography column to determine whether the mixture has an average molecular weight from 4000 to 13,000 Daltons for inclusion in the pharmaceutical product, the improvement comprising

calibrating the molecular weight obtained using the gel permeation chromatography column by subjecting a plurality of molecular weight markers, each of which is a polypeptide consisting essentially of alanine, glutamic acid, tyrosine and lysine and having a predetermined amino acid sequence, to chromatography on the column to establish a relationship between retention time on the column and molecular weight.

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 3 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

124-126. (Canceled)

127. (Previously presented) The process of claim 123, wherein the gel permeation chromatography column comprises a cross-linked agarose-based medium, with an exclusion limit of 2×10^6 Daltons, an optimal separation range of 1000 to 3×10^5 Daltons, and a bead diameter of 20-40 μm .

128. (Previously presented) The process of claim 127, wherein the gel permeation chromatography column is Superose 12.

129. (Previously amended) The process of claim 123, wherein in the molecular weight markers the molar fraction of alanine is 0.38 to 0.5, of glutamic acid is 0.13 to 0.15, of tyrosine is 0.08 to 0.10 and of lysine is 0.3 to 0.4.

130. (Previously Amended) The process of claim 129, wherein in the molecular weight markers the molar fraction of alanine is 0.422 to 0.444, of glutamic acid is 0.133 to 0.143, of tyrosine is 0.086 to 0.093 and of lysine is 0.333 to 0.349.

131. (Previously presented) The process of claim 123, wherein one of the molecular weight markers is selected from the group consisting of

AKKYAKKEKAACKKAYKKEAKAKAAEAAKEAAYEA (SEQ ID NO:1);
AKKYAKKAKAEKAKKAYKAAEAKKAAYEKAAAEKAAKEAAYEA (SEQ ID NO:2);

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 4 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAEAKYKAEAAKAAKEAAAYE
A (SEQ ID NO:3);
AKKYAKKEKAYAKAKKAEAKAAKKAKAEAKKYAKAAKAEKKEYAAAAEAKYKAEAA
KAAAKEAAAYEA (SEQ ID NO:4);
AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKAAAKEAAAYEA (SEQ ID NO:5);
AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKKAYKAEAAKAAAKEAAAYEA (SEQ ID NO:6); and
AKKYAKKAEKAYAKKAKAAKEKKAYAKKEAKAYKAAEAKKKAKAEAKKYAKEAAK
AKKEAYKAEAKKYAKAAKAEKKEYAAAAEAKKAEAAKAYKAEAAKAAAKEAAAYEA
(SEQ ID NO:7).

wherein A represents alanine, K represents lysine, Y
represents tyrosine, and E represents glutamic acid.

132. (Previously presented) The process of claim 123,
wherein the plurality of molecular weight markers is

AKKYAKKEKAACKAYKKEAKAKAAEAAAKEAAAYEA (SEQ ID NO:1);
AKKYAKKAKAEKAKKAYKAAEAKKAAKYEKAAAEKAAAKEAAAYEA (SEQ ID
NO:2);
AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAEAKYKAEAAKAAAKEAAAYE
A (SEQ ID NO:3);
AKKYAKKEKAYAKAKKAEAKAAKKAKAEAKKYAKAAKAEKKEYAAAAEAKYKAEAA
KAAAKEAAAYEA (SEQ ID NO:4);
AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKAAAKEAAAYEA (SEQ ID NO:5);
AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKKAYKAEAAKAAAKEAAAYEA (SEQ ID NO:6); and
AKKYAKKAEKAYAKKAKAAKEKKAYAKKEAKAYKAAEAKKKAKAEAKKYAKEAAK
AKKEAYKAEAKKYAKAAKAEKKEYAAAAEAKKAEAAKAYKAEAAKAAAKEAAAYEA
(SEQ ID NO:7).

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 5 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

wherein A represents alanine, K represents lysine, Y represents tyrosine, and E represents glutamic acid.

133. (Previously Amended) The process of claim 123, wherein the pharmaceutical product is lyophilized.

134. (Previously Amended) A process for obtaining a pharmaceutical product containing a mixture of polypeptides, each of which consists essentially of alanine, glutamic acid, tyrosine and lysine, wherein the mixture has an average molecular weight from 4000 to 13,000 Daltons and in the mixture the molar fraction of alanine is 0.427, of glutamic acid is 0.141, of lysine is 0.337 and of tyrosine is 0.093, which comprises obtaining a batch of a mixture of polypeptides, each of which consists essentially of alanine, glutamic acid, tyrosine and lysine;

determining the average molecular weight of the mixture of polypeptides in the batch using a molecular weight-calibrated gel permeation chromatography column; and

including in the pharmaceutical product the mixture if the mixture is determined to have an average molecular weight from 4000 to 13,000 Daltons,

wherein the gel permeation chromatography column is calibrated by subjecting a plurality of molecular weight markers to chromatography on the column to establish a relationship between the retention time on the column and molecular weight, wherein each of the markers is a polypeptide consisting essentially of

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 6 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

alanine, glutamic acid, tyrosine and lysine and has a predetermined amino sequence.

135-137. (Canceled)

138. (Previously presented) The process of claim 134, wherein the gel permeation chromatography column comprises a cross-linked agarose-based medium, with an exclusion limit of 2×10^6 Daltons, an optimal separation range of 1000 to 3×10^5 Daltons, and a bead diameter of 20-40 μm .

139. (Previously presented) The process of claim 138, wherein the gel permeation chromatography column is Superose 12.

140. (Previously Amended) The process of claim 134, wherein in the molecular weight markers the molar fraction of alanine is 0.38 to 0.5, of glutamic acid is 0.13 to 0.15, of tyrosine is 0.08 to 0.10 and of lysine is 0.3 to 0.4.

141. (Previously Amended) The process of claim 140, wherein in the molecular weight markers the molar fraction of alanine is 0.422 to 0.444, of glutamic acid is 0.133 to 0.143, of tyrosine is 0.086 to 0.093 and of lysine is 0.333 to 0.349.

142. (Previously presented) The process of claim 134, wherein one of the molecular weight markers is selected from the group consisting of

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 7 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

AKKYAKKEKAACKKAYKKEAKAKAAEAAAKEAAAYEA (SEQ ID NO:1);
AKKYAKKAKAEKAKKAYKAAEAKKAAYEKAAAEEKAAAEAAAYEA (SEQ ID
NO:2);
AKKYAKKEKAYAKKAKEKAACKKAEAKAYKAAEAKKKAEAKYKAEAAKAAAEAAAYE
A (SEQ ID NO:3);
AKKYAKKEKAYAKAKKAEAKAAKKAKAEAKKYAKAAKAEKKEYAAAEEKYKAEAA
KAAAEAAAYEA (SEQ ID NO:4);
AKKYAKKEKAYAKKAKEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKAAAEAAAYEA (SEQ ID NO:5);
AKKYAKKEKAYAKKAKEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKAYKAEAAKAAAEAAAYEA (SEQ ID NO:6); and
AKKYAKKAEKAYAKKAKAAKEKKAYAKKEAKAYKAAEAKKKAKAEAKKYAKEAAK
AKKEAYKAEAKKYAKAAKAEKKEYAAAEEKKAEAAKAYKAEAAKAAAEAAAYEA
(SEQ ID NO:7).

wherein A represents alanine, K represents lysine, Y
represents tyrosine, and E represents glutamic acid.

143. (Previously presented) The process of claim 134,
wherein the plurality of molecular weight markers is

AKKYAKKEKAACKKAYKKEAKAKAAEAAAKEAAAYEA (SEQ ID NO:1);
AKKYAKKAKAEKAKKAYKAAEAKKAAYEKAAAEEKAAAEAAAYEA (SEQ ID
NO:2);
AKKYAKKEKAYAKKAKEKAACKKAEAKAYKAAEAKKKAEAKYKAEAAKAAAEAAAYE
A (SEQ ID NO:3);
AKKYAKKEKAYAKAKKAEAKAAKKAKAEAKKYAKAAKAEKKEYAAAEEKYKAEAA
KAAAEAAAYEA (SEQ ID NO:4);
AKKYAKKEKAYAKKAKEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKAAAEAAAYEA (SEQ ID NO:5);
AKKYAKKEKAYAKKAKEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKAYKAEAAKAAAEAAAYEA (SEQ ID NO:6); and

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 8 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

AKKYAKKAEKAYAKKAKAAKEKKAYAKKEAKAYKAAEAKKKAKAEAKKYAKEAAK
AKKEAYKAEAKKYAKAAKAEKKEYAAAEAKKAEAAKAYKAEAAKAAAKEAAAYEA
(SEQ ID NO:7),

wherein A represents alanine, K represents lysine, Y represents tyrosine, and E represents glutamic acid.

144. (Currently Amended) The process of claim 134, further comprising a step of lyophilizing the mixture ~~having the average molecular weight from 4,000 to 13,000 Daltons~~ of polypeptides.

145. (Previously Amended) A process for determining the average molecular weight of an aqueous mixture of polypeptides, each of which consists essentially of alanine, glutamic acid, tyrosine and lysine, wherein in the mixture the molar fraction of alanine is 0.427, of glutamic acid is 0.141, of lysine is 0.337 and of tyrosine is 0.093, which comprises subjecting the mixture to chromatography on a molecular weight-calibrated gel permeation chromatography column so as to determine the average molecular weight of the mixture, wherein the gel permeation chromatography column is calibrated by subjecting a plurality of molecular weight markers to chromatography on the column to establish a relationship between retention time on the column and molecular weight, wherein each of the markers is a polypeptide consisting essentially of alanine, glutamic acid, tyrosine and lysine and has a predetermined amino acid sequence.

146-148. (Canceled)

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 9 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

149. (Previously presented) The process of claim 145, wherein the gel permeation chromatography column comprises a cross-linked agarose-based medium, with an exclusion limit of 2×10^6 Daltons, an optimal separation range of 1000 to 3×10^5 Daltons, and a bead diameter of 20-40 μm .
150. (Previously presented) The process of claim 149, wherein the gel permeation chromatography column is Superose 12.
151. (Previously Amended) The process of claim 145, wherein in the molecular weight markers the molar fraction of alanine is 0.38 to 0.5, of glutamic acid is 0.13 to 0.15, of tyrosine is 0.08 to 0.10 and of lysine is 0.3 to 0.4.
152. (Previously Amended) The process of claim 151, wherein in the molecular weight markers the molar fraction of alanine is 0.422 to 0.444, of glutamic acid is 0.133 to 0.143, of tyrosine is 0.086 to 0.093 and of lysine is 0.333 to 0.349.
153. (Previously presented) The process of claim 145, wherein one of the molecular weight markers is selected from the group consisting of
- AKKYAKKEKAACKKAYKKEAKAKAAEAAAKEAAAYEA (SEQ ID NO:1);
AKKYAKKAKAEKAKKAYKAAEAKKAAYEKA AA EKA AA KEAAAYEA (SEQ ID NO:2);

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 10 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAEAKYKAEAAKAAAKEAAYE
A (SEQ ID NO:3);
AKKYAKKEKAYAKKAEKAACKKAKAEAKKYAKAAKAEKKEYAAAEEAKYKAEAA
KAAAKEAAYEA (SEQ ID NO:4);
AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKAAAKEAAYEA (SEQ ID NO:5);
AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKKAYKAEAAKAAAKEAAYEA (SEQ ID NO:6); and
AKKYAKKAEKAYAKKAKAAKEKKAYAKKEAKAYKAAEAKKKAKAEAKKYAKEAAK
AKKEAYKAEAKKYAKAAKAEKKEYAAAEEAKKAEAAKAYKAEAAKAAAKEAAYEA
(SEQ ID NO:7),

wherein A represents alanine, K represents lysine, Y
represents tyrosine, and E represents glutamic acid.

154. (Previously presented) The process of claim 145,
wherein the plurality of molecular weight markers is

AKKYAKKEKAACKKAYKKEAKAKAAEAAAKEAAYEA (SEQ ID NO:1);
AKKYAKKAKAEKAKKAYKAAEAKKAAYEKAAAEEKAAAKEAAYEA (SEQ ID
NO:2);
AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAEAKYKAEAAKAAAKEAAYE
A (SEQ ID NO:3);
AKKYAKKEKAYAKKAEKAACKKAKAEAKKYAKAAKAEKKEYAAAEEAKYKAEAA
KAAAKEAAYEA (SEQ ID NO:4);
AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKAAAKEAAYEA (SEQ ID NO:5);
AKKYAKKEKAYAKKAEKAACKKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKKAYKAEAAKAAAKEAAYEA (SEQ ID NO:6); and
AKKYAKKAEKAYAKKAKAAKEKKAYAKKEAKAYKAAEAKKKAKAEAKKYAKEAAK
AKKEAYKAEAKKYAKAAKAEKKEYAAAEEAKKAEAAKAYKAEAAKAAAKEAAYEA
(SEQ ID NO:7),

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 11 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

wherein A represents alanine, K represents lysine, Y represents tyrosine, and E represents glutamic acid.

155. (Previously Amended) A process for determining whether an aqueous mixture of polypeptides, each of which consists essentially of alanine, glutamic acid, tyrosine and lysine, has an average molecular weight from 4000 to 13,000 Daltons, wherein in the mixture the molar fraction of alanine is 0.427, of glutamic acid is 0.141, of lysine is 0.337 and of tyrosine is 0.093, which process comprises subjecting the mixture to a calibrated gel permeation chromatography column to determine the average molecular weight of the mixture wherein the gel permeation chromatography column is calibrated by subjecting a plurality of molecular weight markers to chromatography on the column to establish a relationship between retention time on the column and molecular weight, wherein each of the markers is a polypeptide consisting essentially of alanine, glutamic acid, tyrosine and lysine and has a predetermined amino acid sequence.

156-158. (Canceled)

159. (Previously presented) The process of claim 155, wherein the gel permeation chromatography column comprises a cross-linked agarose-based medium, with an exclusion limit of 2×10^6 Daltons, an optimal separation range of 1000 to 3×10^5 Daltons, and a bead diameter of 20-40 μm .

Applicants : Alexander Gad and Dora Lis

Serial No. : 10/792,311

Filed : March 2, 2004

Page 12 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

160. (Previously presented) The process of claim 159, wherein the gel permeation chromatography column is Superose 12.
161. (Previously amended) The process of claim 155, wherein in the molecular weight markers the molar fraction of alanine is 0.38 to 0.5, of glutamic acid is 0.13 to 0.15, of tyrosine is 0.08 to 0.10 and of lysine is 0.3 to 0.4.
162. (Previously amended) The process of claim 161, wherein in the molecular weight markers the molar fraction of alanine is 0.422 to 0.444, of glutamic acid is 0.133 to 0.143, of tyrosine is 0.086 to 0.093 and of lysine is 0.333 to 0.349.
163. (Previously presented) The process of claim 155, wherein one of the molecular weight markers is selected from the group consisting of
- AKKYAKKEKAACKKAYKKEAKAKAAEAAAKEAAYEA (SEQ ID NO:1);
- AKKYAKKAKAEKAKKAYKAAEAKKAACKYEKAAAEKAAAKEAAYEA (SEQ ID NO:2);
- AKKYAKKEKAYAKKAKEAKKAKEAKAYKAAEAKKKAEAKYKAEAAKAAAKEAAYEA (SEQ ID NO:3);
- AKKYAKKEKAYAKKAKEAKKAKEAKKAKAEAKKYAKAAKAEKKEYAAAEAKYKAEAAKAAAKEAAYEA (SEQ ID NO:4);
- AKKYAKKEKAYAKKAKEAKKAKEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA AAEAKYKAEAAKAAAKEAAYEA (SEQ ID NO:5);
- AKKYAKKEKAYAKKAKEAKKAKEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA AAEAKYKAEAAKAYKAEAAKAAAKEAAYEA (SEQ ID NO:6); and

Applicants : Alexander Gad and Dora Lis
Serial No. : 10/792,311
Filed : March 2, 2004
Page 13 of Amendment Under C.F.R. §1.116 in Response to
June 16, 2005 Final Office Action

AKKYAKKAEKAYAKKAKAAKEKKAYAKKEAKAYKAAEAKKKAKAEAKKYAKEAAK
AKKEAYKAEAKKYAKAAKAEKKEYAAAEAKKAEAAKAYKAEAAKAAAKEAAEYEA
(SEQ ID NO:7),

wherein A represents alanine, K represents lysine, Y
represents tyrosine, and E represents glutamic acid.

164. (Previously presented) The process of claim 155,
wherein the plurality of molecular weight markers is

AKKYAKKEKAACKAYKKEAKAKAAEAAAKEAAEYEA (SEQ ID NO:1);
AKKYAKKAKAEKAKKAYKAAEAKKAAKYEKAAAEKAAAKEAAEYEA (SEQ ID
NO:2);
AKKYAKKEKAYAKKAEKAAKAEAKAYKAAEAKKKAEAKYKAEAAKAAAKEAAEYEA
A (SEQ ID NO:3);
AKKYAKKEKAYAKAKKAEAKAAKKAKEAKKYAKAAKAEKKEYAAAEAKYKAEAA
KAAAKEAAEYEA (SEQ ID NO:4);
AKKYAKKEKAYAKKAEKAAKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKAAAKEAAEYEA (SEQ ID NO:5);
AKKYAKKEKAYAKKAEKAAKAEAKAYKAAEAKKKAKAEAKKYAKAAKAEKKEYA
AAEAKYKAEAAKAYKAEAAKAAAKEAAEYEA (SEQ ID NO:6); and
AKKYAKKAEKAYAKKAKAAKEKKAYAKKEAKAYKAAEAKKKAKAEAKKYAKEAAK
AKKEAYKAEAKKYAKAAKAEKKEYAAAEAKKAEAAKAYKAEAAKAAAKEAAEYEA
(SEQ ID NO:7),

wherein A represents alanine, K represents lysine, Y
represents tyrosine, and E represents glutamic acid.